## High Latitude Survey Figure of Merit

(HLS FoM)

C. Conselice, P. Schechter, D. Stern, E. Wright

#### Outline

- 1. High Redshift QSOS (DS)
- 2. Galaxy Evolution/Structure (CC)
- 3.Cool and rare stars (EW)

# I. High-Redshift Quasars

- We define two QSO FoM's, based on two distinct scientific questions:
  - The small number of brightest, highest redshift QSOs, which are the best background probes of the intervening universe
  - The large number of z>6 QSOs, going all the way down the QLF: faintend of the QLF, AGN contribution to ionizing budget, QSO pairs, LSS, clustering to get halo masses, etc...
- Define FOMQ1 as the normalized number of QSOs detected by WFIRST at z>8 with M(1450) < -25 (corresponding to what SDSS does at  $z\sim4$ ):

$$FOM_{Q1} = \kappa \int \int \Phi (M_{1450}, z)$$

where K is the normalization, the integrals are over the above specified ranges of absolute magnitude and redshift.


QSO FoM #2: The large number of QSOs at z>6.

### Normal Galaxies FoM

Number of galaxies, N or 10^N or completeness depth of L\*+N(mag)

FoM – fraction of galaxies resolved to some magnitude limit, i.e., H = 25, real example from Hubble UDF (right) finds FoM = 0.77 for 0.25 arcsec PSF

Drop-outs: ultra-high redshift galaxies, z > 7: (to depth of H~26)




FoM (z-drops) = 610\*sq. degree surveyed

FoM (Y-drops) = 90\*sq. degree surveyed

FoM (J-drops) = 80\*sq. degree surveyed

#### Rare Stars

- Searching for rare stars requires a wide range of filters to separate them from the MS locus.
- WD spectrum is suppressed at 1.1 and 2.2 µm so optical and YJHKL filters are good.
- BD spectrum peaks at JH and M, so filters from 1-3.5 μm are good, 4.6 μm is ideal.
- Hundreds of cool WDs and cold BDs can be found.
- Euclidean source counts strongly favor wide survey.



- 4000 K WD showing H<sub>2</sub> CIA
- 325 K BD showing CH<sub>4</sub> dominated spectrum
- Best MS match to WD